ZU-TH 12/95
hep-th/9609201

arXiv:hep-th/9609201v1 25 Sep 1996

Negative modes around Einstein–Yang–Mills sphalerons and black holes[1]

Mikhail S. Volkov

Institut für Theoretische Physik der Universität Zürich–Irchel,
Winterthurerstrasse 190, CH–8057 Zürich, Switzerland,
e–mail: volkov@physik.unizh.ch

Abstract

The dynamics of small perturbations around sphaleron and black hole solutions in the Einstein-Yang-Mills theory for the gauge group $SU(2)$ is investigated. The perturbations can be split into the two independent sectors in accordance with their parity; each sector contains negative modes. The even-parity negative modes are shown to correspond to the negative second variations of the height of the potential energy barrier near the sphaleron. For the odd-parity sector, the existence of precisely n (the number of nodes of the sphaleron solution gauge field function) negative modes is rigorously proven. The same results hold for the Einstein-Yang-Mills black holes as well.

---

[1] Published in *Geometry and Integrable Models*, Eds. P.N. Pyatov & S.N.Solodukhin, World Scientific 1996, pp. 55-77.

1 Introduction

Soon after the discovery of the Bartnik-McKinnon solitons [1] and black holes [2] in the $SU(2)$ Einstein-Yang-Mills (EYM) theory, it was shown that these interesting objects are unfortunately unstable [3]. However, as has been realized later [4], the property to be unstable is in fact fairly important in the case, since it allows for the sphaleron interpretation [5],[6] of the regular BK objects.

Usually, sphalerons are related with the existence of non-contractible loops in the configuration space of the theory [5], [7]. For each such a loop, one finds a maximal value of the potential energy and then minimizes the result over all loops; this gives the energy of a static saddle point field configuration. Notice that the configuration space has to be compact in this approach, otherwise the existence of non-contractible loops does not ensure the existence of saddle points of the energy [5]. The difference between the two cases is perfectly illustrated in the two figures in famous Manton's paper (see Fig.1 and Fig.2 in [5]). Although for the non-compact case the saddle points may also exist (see Fig.2 in [5]), these points have *more then one* directions of instability, and thus can not be constructed in the same manner is in the compact case, such that their existence is not directly related to the existence of non-contractible loops. The use of loops in the non-compact case may have another meaning: for theories with vacuum periodicity, the non-contractible loop through the saddle point (if any exists) can be transformed into a path interpolating between distinct vacua. This shows that the saddle point field configuration relates to the top of the potential barrier between the vacua thus justifying the usage of the name "sphaleron".

It is not widely understood that the EYM sphalerons correspond to the "non-compact" case. In the EYM theory, the maximal value of energy for each non-contractible loop can be reduced to zero by a smooth deformation of the loop [4], such that the existence of the loops *does not ensure* the existence of the solutions (otherwise the existence proof for the BK solutions [8] – [10] would just follow the same line as that for the standard sphaleron [11]). Instead, in order to illustrate the sphaleron nature of the solutions, the loops can be used as paths connecting the solutions with the neighboring vacua [12] (see below). Manifestly, the difference between the standard Yang-Mills-Higgs (YMH) sphaleron and the EYM sphalerons results in the different numbers of their unstable modes. The YMH sphaleron [5] has one and only one negative mode in the odd-parity spherically-symmetric perturbation sector. In the even-parity sector, the solution has no negative modes, such that the energy of the solution indeed defines the *minimal* height of the barrier. The EYM sphalerons have negative modes both in the odd parity and in the even-parity sectors, which means that they do not specify the minimal possible height of the barrier (the latter is zero) [13].

The present paper concerns with the investigation of the dynamics of spherically-symmetric perturbation modes around EYM sphalerons and black holes. The perturbations decouple into the two independent groups in accordance with their parity. The even parity modes were studied previously in [3], it has been found numerically that the n-th soliton or black hole solution has precisely n negative modes of this type. Here we explicitly show that the negative modes of this type are related to the decrease of the *height* of the potential barrier around the sphaleron [13]. For the odd-parity modes, the existence of at least one such a mode for all known $SU(2)$ EYM solitons and black holes has been established in [4], [14], as well as for the "generic" solitons in the EYM theory for any compact gauge group [15]. For the lowest n regular BK solutions, numerical analysis has revealed the existence of n negative modes of this type too [16]. Here we give a complete proof of the following statement: for all known regular and black hole non-Abelian EYM solutions there exist precisely n negative modes in the odd parity spherically-symmetric perturbation sector [17]. For the even-parity modes the corresponding proof is unknown, however the results of the numerical analysis suggest that there are n negative modes in this case too, such that the total number of negative modes for the EYM solitons and black holes is $2n$.

The rest of the paper is organized as follows. In Sec.2 the basic notations and equations are introduced. Sec.3 contains an explicit construction of vacuum-to-vacuum paths through the EYM sphalerons. In Sec.4 the basic perturbation equations are given for the both parity types. For the odd-parity perturbation modes, the transformation to a one-channel Schrödinger problem is given in Sec.5, the regularity of the effective potential is shown in Sec.6. A zero-energy bound state in the problem is explicitly presented in Sec.7, the corresponding wave function has n zeros, which proves the existence of precisely n bound states. The black hole case is considered in Sec.8, the results apply also for the case of charged non-Abelian black holes found in the EYM theory with the $SU(2)\times U(1)$ gauge group [18]. The second variation of the ADM mass functional for the regular BK solutions is explicitly derived in the Appendix.

2 EYM field equations

We start from the action of the $SU(2)$ EYM theory

$$S = -\frac{1}{16\pi G}\int R\sqrt{-g}d^4x - \frac{1}{2e^2}\int tr F_{\mu\nu}F^{\mu\nu}\sqrt{-g}d^4x, \tag{1}$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]$ is the matrix valued gauge field tensor, e is the gauge coupling constant, $A_\mu = A_\mu^a\tau^a/2$, and τ^a $(a = 1, 2, 3)$ are the Pauli matrices.

In the spherically symmetric case, it is convenient to represent the spacetime metric in the following form:

$$ds^2 = l_e^2\left((1-\frac{2m}{r})\sigma^2 dt^2 - \frac{dr^2}{1-2m/r} - r^2(d\vartheta^2 + \sin^2\vartheta d\varphi^2)\right), \tag{2}$$

where m and σ depend on t and r. Here the EYM length scale $l_e = \sqrt{4\pi G}/e$ is explicitly displayed, after such a rescaling, all other quantities in the theory become dimensionless.

The spherically symmetric $SU(2)$ Yang-Mills field can be parameterized by

$$A = W_0\hat{L}_1\, dt + W_1\hat{L}_1\, dr + \{p_2\,\hat{L}_2 - (1-p_1)\,\hat{L}_3\}\, d\vartheta + \{(1-p_1)\,\hat{L}_2 + p_2\,\hat{L}_3\}\sin\vartheta\, d\varphi, \tag{3}$$

where W_0, W_1, p_1, p_2 are functions of t and r, $\hat{L}_1 = n^a\tau^a/2$, $\hat{L}_2 = \partial_\vartheta\hat{L}_1$, $\sin\vartheta\hat{L}_3 = \partial_\varphi\hat{L}_1$, and $n^a = (\sin\vartheta\cos\varphi, \sin\vartheta\sin\varphi, \cos\vartheta)$. The gauge transformation

$$A \to\; UAU^{-1} + iUdU^{-1}, \;\; \text{with}\;\; U = \exp(i\Omega(t,r)\hat{L}_1), \tag{4}$$

preserves the form of the field (3), altering the functions W_0, W_1, p_1, p_2 as

$$W_0 \to W_0 + \partial_t\Omega, \quad W_1 \to W_1 + \partial_r\Omega, \quad p_\pm = p_1 \pm ip_2 \to \exp(\pm i\Omega)p_\pm. \tag{5}$$

Consider also parity transformation: $\vartheta \to \pi - \vartheta$, $\varphi \to \varphi + \pi$. The action of this transformation on the fields (2), (3) is equivalent to the following replacement:

$$m \to m, \quad \sigma \to \sigma, \quad p_1 \to p_1, \quad p_2 \to -p_2, \quad W_0 \to -W_0, \quad W_1 \to -W_1. \tag{6}$$

It is convenient to introduce the complex variable $f = p_1 + ip_2$ and its covariant derivative $D_\mu f = (\partial_\mu - iW_\mu)f$, as well as the $(1+1)$-dimensional field strength $W_{\mu\nu} = \partial_\mu W_\nu - \partial_\nu W_\mu$, $(\mu, \nu = 0, 1)$. The full system of the EYM equations then reads

$$\partial_\mu(r^2\sigma W^{\mu\nu}) - 2\sigma \; Im \; (fD^\nu f)^* = 0, \tag{7}$$

$$D_\mu \sigma D^\mu f - \frac{\sigma}{r^2}(|f|^2 - 1)f = 0, \tag{8}$$

$$\partial_r m = -\frac{r^2}{4}W_{\mu\nu}W^{\mu\nu} + \frac{1}{N\sigma^2}|D_0 f|^2 + N|D_1 f|^2 + \frac{1}{2r^2}(|f|^2-1)^2, \tag{9}$$

$$\partial_t m = 2N \; Re \; D_0 f (D_1 f)^*, \tag{10}$$

$$\partial_r \ln \sigma = \frac{2}{r}\left(\frac{1}{N^2\sigma^2}|D_0 f|^2 + |D_1 f|^2\right), \tag{11}$$

where $N = 1 - 2m/r$, asterisk denotes complex conjugation.

In the static case, with $W_0 = W_1 = p_2 = 0$ and $f = Re \; f = p_1$, these equations are known to possess regular, asymptotically flat solutions discovered by Bartnik and McKinnon (BK) [1], as well as the EYM black hole solutions [2]. The regular solutions form a discrete family labeled by an integer $n = 1, 2, \ldots$. Near the origin and infinity, these solutions, $(f_n(r), m_n(r)\sigma_n(r))$, have the following common behaviour:

$$f = 1 - br^2 + O(r^4), \quad m = 2b^2r^3 + O(r^5), \quad \frac{\sigma}{\sigma_0} = 1 + 4b^2r^2 + O(r^4), \quad \text{as} \quad r \to 0, \tag{12}$$

and

$$f = (-1)^n(1 - \frac{a}{r} + O(\frac{1}{r^2})), \quad m = M + O(\frac{1}{r^3}), \quad \sigma = 1 + O(\frac{1}{r^4}) \;\; \text{as} \;\; r \to \infty, \tag{13}$$

where $0 < b, \sigma_0 < 1$, M and a are numerically known constants depending on n. In the domain $0 < r < \infty$, $m(r)$ and $\sigma(r)$ are monotone increasing functions, while $f(r)$ oscillates n times around zero value always staying within the stripe $-1 < f(r) < 1$.

The EYM black hole solutions are distinguished by the two parameters, n and r_h, where n is the number of nodes of the function $f(r)$ in the region $r > r_h$, and $r_h \in (0, \infty)$ is the event horizon size. The boundary conditions at the horizon $r = r_h$ are

$$f = f_h + \frac{P}{F}\,x + O(x^2), \quad N = F\,x + O(x^2), \quad \frac{\sigma}{\sigma_h} = 1 + \frac{2P^2}{r_h^2F^2}\,x + O(x^2), \tag{14}$$

where

$$x = \frac{r - r_h}{r_h}, \quad P = f_h(f_h^2 - 1), \quad F = 1 - (f_h^2 - 1)^2/r_h^2, \tag{15}$$

and $0 < f_h, \sigma_h$ depend on (n, r_h). In the domain $r_h < r < \infty$ the behaviour of these solutions is qualitatively similar to that for the regular case with the boundary conditions at infinity defined by (13).

In what follows, it will be convenient to introduce the tortoise coordinate ρ such that $d\rho/dr = 1/\sigma N$. In the regular case ρ runs over the half-line $[0, \infty)$:

$$\rho = \frac{r}{\sigma_0} + O(r^5) \ \text{ as } \ r \to 0, \qquad \rho = r + 2M \ln r + O(1) \ \text{ as } \ r \to \infty. \tag{16}$$

For the black hole solutions the domain of ρ is the whole line $(-\infty, +\infty)$ such that near the horizon

$$N = 1 - \frac{2m}{r} = F \exp\left(\sigma_h F \frac{\rho}{r_h}\right) + O\left(\exp\left(2\sigma_h F \frac{\rho}{r_h}\right)\right) \ \text{ as } \ \rho \to -\infty. \tag{17}$$

3 EYM sphalerons and vacuum to vacuum paths

As was shown in Ref.[4], the odd-n BK solutions may be treated as sphalerons. This interpretation is based on the fact that, through any such a solution, one may find a one-parameter family of static field configurations which interpolates between topologically distinct EYM vacua. If λ denotes the parameter of the family, the corresponding gauge field can be chosen as follows [4]

$$A_\mu dx^\mu = \frac{i}{2e}(1 - f_n(r)) U dU^{-1}, \quad U = exp(i\lambda n^a \tau^a), \tag{18}$$

where λ ranges from zero to π. The metric functions m and σ in (2) are given by

$$\sigma(r) = exp\{-2sin^2\lambda \int_r^\infty f_n'^2 \frac{dr}{r}\},$$

$$m(r) = \frac{sin^2\lambda}{\sigma(r)} \int_0^r (f_n'^2 + sin^2\lambda \frac{(f_n^2 - 1)^2}{2r^2}) \sigma dr. \tag{19}$$

It is worth noting that, for any value of λ, the fields of this family satisfy the two Einstein equations $G_0^0 = 8\pi G T_0^0$ and $G_r^r = 8\pi G T_r^r$; when $\lambda = \pi/2$ these equations reduce to Eqs.(9), (11) [4]. One can check that, when $\lambda = \pi/2$, the fields defined by (18), (19) coincide with the n-th BK solution field described above. When λ approaches zero or π values, the spacetime metric becomes flat and the gauge field (18) vanishes. Thus, the family of fields (18), (19) describes a loop (in the EYM configuration function space) which interpolates between the trivial vacuum and the n-th BK solution when λ runs from 0 to $\pi/2$, and returns back to the vacuum as λ changes from $\pi/2$ to π.

It is important that this loop can be transformed to a path interpolating between neighboring EYM vacua with different winding numbers of the gauge field. To see this, one has to pass to such a gauge, in which the gauge field potential decays faster then $1/r$ as $r \to \infty$. The gauge transformation

$$A_\mu \to U(A_\mu + \frac{i}{e}\partial_\mu)U^{-1}, \ \text{ with } \ U = exp(i\frac{\lambda}{2}(f_n - 1)n^a \tau^a) \tag{20}$$

yields

$$A_\mu dx^\mu = \frac{i}{2e}(1 - f_n(r)) U_+ \, dU_+^{-1} + \frac{i}{2e}(1 + f_n(r)) U_- \, dU_-^{-1},$$

$$U_\pm = exp(i\lambda(f_n \pm 1)n^a \tau^a/2), \tag{21}$$

whereas the gauge invariant metric functions (19) remain unchanged. One can see that, when λ runs from zero to π, the field (21) interpolates between two different pure gauges. Notice that, in the new gauge, one has $A_a = O(r)$ when $r \to 0$ and $A_a = O(1/r^2)$ as $r \to \infty$, with A_a being the field component with respect to an orthonormal frame.

Let λ depend adiabatically on time, in such a way that $\lambda(t = -\infty) = 0$ and $\lambda(t = \infty) = \pi$. Notice that the structure of the field (21) ensures the temporal gauge condition $A_0 = 0$ even for time-dependent λ. Recall the Chern-Simons (CS) current and its divergence

$$K^\mu = \frac{e^2}{8\pi^2} tr \frac{\varepsilon^{\mu\nu\alpha\beta}}{\sqrt{-g}} A_\nu (\nabla_\alpha A_\beta - \frac{2ie}{3} A_\alpha A_\beta), \quad \nabla_\alpha K^\alpha = \frac{e^2}{16\pi^2} tr F_{\mu\nu} \tilde{F}^{\mu\nu}, \tag{22}$$

where ∇_α is the covariant derivative, and $\tilde{F}^{\mu\nu}$ is dual tensor. Integration leads to the following relation:

$$\frac{e^2}{16\pi^2} \int_{-\infty}^{t} dt \int d^3x \sqrt{-g} tr F_{\mu\nu} \tilde{F}^{\mu\nu} = \int d^3x \sqrt{-g} K^0 \Big|_{-\infty}^{t} + \int_{-\infty}^{t} dt \oint \vec{K} d\vec{\Sigma}. \tag{23}$$

In the temporal gauge used, the field decays faster then $1/r$ as $r \to \infty$, such that the surface integral entering the right hand side vanishes, while the gauge invariant left hand side is [4]

$$\frac{3}{2\pi} \int_{-\infty}^{t} dt\ \dot{\lambda} sin^2\lambda \int_0^\infty dr\ f_n'(f_n^2 - 1) = \frac{3}{4\pi}(\lambda - sin\lambda cos\lambda)(\frac{1}{3} f_n^3 - f_n)\Big|_{f_n(0)=1}^{f_n(\infty)=-1}. \tag{24}$$

Noting that the gauge potential (21) vanishes at $\lambda = 0$, one obtains the following value of the CS number along the path (21) [4]:

$$N_{CS} = \int \sqrt{-g} K^0 d^3x \Big|_t = \frac{1}{\pi}(\lambda - sin\lambda cos\lambda), \tag{25}$$

which changes from zero to one as λ runs from 0 to π.

Thus the fields (2), (19), (21) interpolate between two EYM vacua with different winding numbers of the gauge field, and coincide with the (gauge transformed) field of the n-th BK solution when $\lambda = \pi/2$. This allows one to treat odd-n BK solutions as sphalerons, “lying” on the top of the potential barrier separating distinct topological vacuum sectors of the theory. The profile of this barrier may be obtained from Eq.(19) due to the fact that the metric function $m(r)$ obeys the initial value constraint, which allows one to define the ADM energy [4] [4], [13]

$$E[\lambda, f(r)] = \lim_{r\to\infty} m(r) =$$
$$= sin^2\lambda \int_0^\infty (f'^2 + sin^2\lambda \frac{(f^2-1)^2}{2r^2}) exp(-2 sin^2\lambda \int_r^\infty f'^2 \frac{dr}{r}) dr, \tag{26}$$

such that the sphaleron energy is given by $E[\pi/2, f_n(r)]$.

4 Perturbations of the equilibrium solutions

The functional E given by (26) is quite useful for a qualitative understanding of the behaviour of the potential barrier surface in the vicinity of sphalerons [13]. The sphaleron configurations, $\lambda = \pi/2, f(r) = f_n(r)$, correspond to critical points of $E[\lambda, f(r)]$. One can easily see that $E[\lambda, f(r)]$ reaches its maximal value at $\lambda = \pi/2$, which shows that each sphaleron possesses an unstable mode [4]; this kind of instability can be naturally called sphaleron instability. In addition, the value of $E[\pi/2, f_n(n)]$ decreases under certain deformations of another type: $f_n \to f_n + \delta f$ [13] which will be referred to as gravitational instabilities.

To put the this into a more rigorous form, we return to the general system of the EYM field equations (7)-(11) and consider small spherically-symmetric perturbations of a given (regular or black hole) equilibrium solution:

$$m \to m + \delta m, \quad \sigma \to \sigma + \delta\sigma, \quad f \to f + \delta f, \quad W_0 = W_1 = 0 \to \delta W_0, \; \delta W_1. \tag{27}$$

In view of (6), these perturbations can be classified in accordance with their parity. Observing that the background equilibrium solutions are invariant under parity, one concludes that even–parity perturbations, $(\delta m, \delta\sigma, \delta p_1)$, and odd–parity perturbations, $(\delta W_0, \delta W_1, \delta p_2)$, must be independent, which is confirmed by the straightforward linearization of Eqs.(7)-(11). Notice that the infinitesimal gauge transformation (5) does not alter the even-parity perturbations, while the odd-parity ones change as

$$\delta W_0 \to \delta W_0 + \partial_t \Omega, \quad \delta W_1 \to \delta W_1 + \partial_r \Omega, \quad \delta p_2 \to \delta p_2 + \Omega f. \tag{28}$$

Perturbations in the even-parity sectors have been studied in [3]. In this case, linearization of Eqs.(8), (10), (11) reveals that perturbations of the metric functions, δm and $\delta\sigma$, can be expressed entirely in terms of δp_1, the latter satisfies the following Schrödinger-type equation:

$$-\phi'' + \left(\sigma^2 N \frac{3f^2 - 1}{r^2} + 2\left(\frac{\sigma'}{\sigma}\right)'\right)\phi = \omega^2 \phi, \tag{29}$$

where $\delta p_1 = \exp(-i\omega t)\phi(\rho)$ with ρ being the tortoise coordinate introduced above (throughout this paper we use the following notations: $y' = dy/d\rho$, and $y'_r = dy/dr$). For the n-th background regular or black hole solution, numerical analysis has shown the existence of n bound states in (29).

The even parity perturbations correspond to the gravitational perturbations defined above. It is instructive to see how the perturbation equation (29) can be derived directly from the potential barrier functional E [13]. Let us introduce *a barrier height functional*

$$\varepsilon[f(r)] = E[\lambda = \pi/2, f(r)], \tag{30}$$

and consider its expansion near a critical point

$$\varepsilon[f_n(r) + \phi(r)] = \varepsilon[f_n(r)] + \delta^2\varepsilon + \ldots, \tag{31}$$

where the first order term vanishes and dots denote higher order terms. The second order term reads

$$\delta^2\varepsilon = \int_0^\infty \phi(-\frac{d^2}{d\rho^2} + V)\phi d\rho, \tag{32}$$

The effective potential V in this expression is explicitly derived in the Appendix, and turns out to be the same as that in the Schrödinger equation (29). This means that any solution of the perturbation problem (29) with $\omega^2 < 0$ gives rise to the negative second variation of the barrier height functional.

Let us pass now to the analysis of the odd-parity modes. The odd-parity perturbation sector contains the sphaleron instabilities of the equilibrium solutions. For the odd-parity perturbations, the metric remains unperturbed, such that the perturbation equations can easily be obtained via expanding the Yang-Mills equations (7), (8) with respect to δW_0, δW_1, and δp_2 alone. We use the gauge freedom (28) to impose the condition $\delta W_0 = 0$, and specify the time dependence as $\delta W_1 = \exp(-i\omega t)\alpha(r)$, $\delta p_2 = \exp(-i\omega t)\xi(r)$. The resulting equations read

$$\left(-\frac{d}{dr}N\sigma\frac{d}{dr} + \frac{\sigma}{r^2}(f^2-1)\right)\xi + (f'_r + \frac{d}{dr}f)N\sigma\alpha = \frac{\omega^2}{\sigma N}\xi, \tag{33}$$

$$2N\frac{\sigma^2}{r^2}\left(f^2\alpha + f'_r\xi - f\xi'_r\right) = \omega^2\alpha, \tag{34}$$

where f, σ and N refer to the background equilibrium solution. There exists also an additional equation due to the Gauss constraint (Eq.(7) with $\nu = 0$):

$$\omega(\left(\frac{r^2}{\sigma}\alpha\right)'_r - \frac{2f}{N\sigma}\xi) = 0. \tag{35}$$

To simplify these equations we introduce the new variable $\chi = (r^2/2\sigma)\alpha$ and pass to the tortoise coordinate ρ. Multiplying (33) by f and using the background Yang-Mills equation

$$f'' = \frac{\sigma^2 N}{r^2}f(f^2-1), \tag{36}$$

one can represent Eqs.(33-35) in the following form:

$$\left(f^2\gamma^2\chi + f'\xi - f\xi'\right)' = \omega^2 f\xi, \tag{37}$$

$$f^2\gamma^2\chi + f'\xi - f\xi' = \omega^2\chi, \tag{38}$$

$$\omega(\chi' - f\xi) = 0, \tag{39}$$

where $\gamma^2 = 2N\sigma^2/r^2$. One can immediately see that, as long as $\omega \neq 0$, only two of these equations are independent, say, (38) and (39) (one can equally say that Eqs.(37), (38) are independent, and the Gauss constraint in the case is the differential consequence of this equation of motion). When $\omega = 0$, Eq.(39) disappears, and there remains only one essential equations (38) which can be solved by any static pure gauge obtained from (28):

$$\chi = \frac{\Omega'}{\gamma^2}, \quad \xi = f\Omega. \tag{40}$$

Let ω be non-vanishing. Then one can omit the factor ω in (39) and specify the following independent equations which are clearly equivalent to (37)-(39):

$$f\xi' - f'\xi = (f^2\gamma^2 - \omega^2)\chi, \tag{41}$$

$$\chi' = f\xi. \tag{42}$$

When $\omega = 0$, Eq.(42) does not appear in the initial system, however we will retain this equation at zero energy too. In this case, this equation (it is sometimes called "strong

Gauss constraint") plays the role of the gauge fixing condition. Imposing this condition violates the remaining gauge invariance of equations (37)-(39) at zero energy, allowing only for those pure gauge solutions (40) whose parameter Ω satisfies

$$\left(\frac{\Omega'}{\gamma^2}\right)' = f^2\Omega. \tag{43}$$

5 Transformation to Schrödinger problem

Our aim now is to reduce Eqs.(41), (42) to a one-channel Schrödinger problem. The most natural way to do this is to exclude one of the functions, either ξ or χ, from the equations. In this case however, one obtains either singular or energy-dependent effective potential in the resulting Schrödinger equation. For instance, eliminating ξ one finds

$$-\phi'' + \left(\gamma^2(f^2+1) + 2\left(\frac{f'}{f}\right)^2\right)\phi = \omega^2\phi, \tag{44}$$

where $\phi = \chi/f$, the potential in this equation is ill-defined due to nodes of f. Another possibility is to try to look for a certain linear combination of χ and ξ which might satisfy a Schrödinger-type equation with regular, energy-independent potential. Surprisingly, this way turns out to be successful. The key observation here is that the combination

$$\psi = fZ\chi + \xi, \tag{45}$$

where Z is a solution of the following auxiliary equation:

$$Z' = f^2Z^2 - \gamma^2, \tag{46}$$

satisfies by virtue of (41), (42) and (46) the Schrödinger-type equation

$$-\psi'' + U\psi = \omega^2\psi, \tag{47}$$

with the energy-independent effective potential

$$U = \sigma^2 N\frac{3f^2-1}{r^2} + 2\left(f^2Z\right)'. \tag{48}$$

The inverse transformation from ψ to the variables χ and ξ reads

$$\chi = \frac{1}{\omega^2}\left(f^3Z\psi + f'\psi - f\psi'\right), \qquad \xi = \psi - fZ\chi. \tag{49}$$

In the new potential derived the nodes of f are no more dangerous, such that the potential is regular provided that Z is regular. Furthermore, the investigation of the behaviour of Z is simplified considerably due the existence of another remarkable relation allowing us to represent solutions of Eq.(46) in terms of solutions of second-order *linear* differential equation (43):

$$Z = -\Omega\Lambda - \frac{\Lambda^2}{C + \int_0^\rho f^2\Lambda^2 d\rho}, \qquad \text{where} \quad \Lambda = \frac{\gamma^2}{\Omega'}, \tag{50}$$

with C being an integration constant. For each particular $\Omega(\rho)$, this correspondence provides us with a family of solutions to Eq.(46) whose members are distinguished by values of C in (50). However, not all of these solutions are well-behaved. Our aim is to specify Ω and C in (50) such that the resulting $Z(\rho)$ gives rise to well-defined potential (48) and transformations (45), (49).

Before passing to the further analysis, one may wonder why it was possible to guess the transformation presented. One might suspect the existence of some symmetry in the problem, which could naturally explain the appearance of all the relations listed above. It turns out that there exists indeed a regular way to proceed in the case, since the transformation applied formally resembles the transformation between the dual partners in supersymmetric quantum mechanics [17]. First, notice that Eq.(44) has obvious $\omega = 0$ solutions which are just pure gauge modes (see (40)),

$$\phi_0 = \Omega'/f\gamma^2, \tag{51}$$

with Ω satisfying (43). Any such solution allows us to factorize the differential operator in (44):

$$-\frac{d^2}{d\rho^2} + \left(\gamma^2(f^2+1) + 2\left(\frac{f'}{f}\right)^2\right) = -\frac{d^2}{d\rho^2} + \frac{\phi_0''}{\phi_0} = Q^+Q^-, \tag{52}$$

with

$$Q^\pm = \mp\frac{d}{d\rho} - \frac{\phi_0'}{\phi_0}. \tag{53}$$

Using (43) we find

$$Q^\pm = \mp\frac{d}{d\rho} - \frac{f'}{f} - f^2 Z, \tag{54}$$

where

$$Z = -\Omega\Lambda, \quad \Lambda = \gamma^2/\Omega'. \tag{55}$$

Notice that (43) is equivalent to

$$\Lambda'/\Lambda = f^2 Z, \tag{56}$$

and thus Z satisfies the nonlinear differential equation (46).

With a standard reduction, the most general solution of the second order linear equation (43) is

$$\Omega = c_2\tilde{\Lambda} + \tilde{\Omega}\left(c_1 + c_2\int_0^\rho f^2\tilde{\Lambda}^2 d\rho\right), \tag{57}$$

where $\tilde{\Omega}$ is a special solution, $\tilde{\Lambda} = \gamma^2/\tilde{\Omega}'$, and c_1, c_2 are real constants. This gives immediately for Z in (55)

$$Z = -\tilde{\Omega}\tilde{\Lambda} - \frac{\tilde{\Lambda}^2}{c_1/c_2 + \int_0^\rho f^2\tilde{\Lambda}^2 d\rho}, \tag{58}$$

which is obviously identical to (50). Since the differential equation (44) is, for any Ω in (57), identical to

$$Q^+Q^-\phi = \omega^2\phi, \tag{59}$$

one can pass from ϕ to

$$\psi = Q^-\phi. \tag{60}$$

For $\omega \neq 0$ this has the unique inverse

$$\phi = \frac{1}{\omega^2} Q^+ \psi \tag{61}$$

and by applying Q^- we obtain the "dual" eigenvalue equation

$$Q^- Q^+ \psi = \omega^2 \psi. \tag{62}$$

The differential operator on the left is

$$Q^- Q^+ = -\frac{d^2}{d\rho^2} + U, \tag{63}$$

with U being given by (48), whereas transformations (60), (61) are identical to (45), (49).

6 Structure of the potential

Now we turn to the investigation of the behaviour of Z. First, consider the case of the regular background equilibrium solutions. It is convenient to return for a moment to the variable r, then Eq.(43) takes the form

$$r^2 \Omega''_{rr} + 2r(1 - f'^2_r)\Omega'_r = \frac{2f^2}{N}\Omega. \tag{64}$$

This differential equation has regular singular points at $r = 0, \infty$. As is known [10], the background solutions (12), (13) are analytic in some neighborhoods of the origin and infinity thus ensuring analyticity of the coefficients in (64). This guarantees the existence of the fundamental system of solutions in the vicinity of each singular point [19]. Using (12), (13), (64) one can represent these solutions in the following form:

$$\Omega(r) = c_1 \; r \left\{1 + R_1(r)\right\} + c_2 \; \frac{1}{r^2} \left\{1 + R_2(r)\right\} \qquad \text{as} \;\; r \to \;\; 0, \tag{65}$$

and introducing $s = 1/r$,

$$\Omega(s) = d_1 \; s^2 \left\{1 + S_1(s)\right\} + d_2 \; \frac{1}{s} \Big\{1 + S_2(s) + \mathcal{A} \; s^3 ln(s)(1 + S_1(s))\Big\} \qquad \text{as} \;\; s \to \;\; 0, \tag{66}$$

where $c_{1,2}$ and $d_{1,2}$ are arbitrary constants, $R_{1,2}(r)$ and $S_{1,2}(s)$ are convergent power series vanishing at $r = 0$ and $s = 0$, respectively, the quantity $\mathcal{A} = \mathcal{A}(a, M)$ is specified by the asymptotic behaviour of the background solution.

In the domain $0 < r < \infty$ Eq.(64) has no singular points and the coefficients of the equation are at least continuous functions in this region. This guarantees the existence of an extension of the asymptotic solutions (65), (66) to the whole half-line, and this extension is at least C^2.

Now we choose a special solution $\tilde{\Omega}$ which is regular at infinity. For this we specify in (66) $d_1 = 1$, $d_2 = 0$. Then, near the origin, $\tilde{\Omega}$ is given by (65). One can see that the coefficient c_2 in this formula does not vanish. Indeed, multiplying Eq.(43) by Ω and integrating from ρ to infinity one obtains

$$0 = \int_\rho^\infty (-\Omega \left(\frac{\Omega'}{\gamma^2}\right)' + f^2\Omega^2) d\rho = -\frac{\Omega'}{\gamma^2}\Omega\Big|_\rho^\infty + \int_\rho^\infty (\frac{\Omega'^2}{\gamma^2} + f^2\Omega^2) d\rho. \tag{67}$$

Applying this relation to the particular solution $\tilde{\Omega}$, $\tilde{\Omega}(\infty) = \tilde{\Omega}'(\infty) = 0$, one arrives at

$$(\tilde{\Omega}^2)' = -2\gamma^2 \int_\rho^\infty (\frac{\tilde{\Omega}'^2}{\gamma^2} + f^2\tilde{\Omega}^2) d\rho < 0, \tag{68}$$

so that, since $\tilde{\Omega}$ vanishes at infinity, it can not vanish also at the origin containing therefore the growing branch, $\tilde{\Omega} \sim 1/r^2$ as $r \to 0$. From here we conclude that on the domain $0 < r < \infty$ there exists a smooth (at least C^2) solution $\tilde{\Omega}(r)$ with the following behaviour near $r = 0, \infty$:

$$\tilde{\Omega} = \frac{const}{r^2} + O(1) \ \text{ as } r \to 0, \qquad \tilde{\Omega} = \frac{1}{r^2} + O(\frac{1}{r^3}) \ \text{ as } r \to \infty. \tag{69}$$

Next, we substitute this solution in Eq.(50), where we require the constant C to be positive – to avoid vanishing of the denominator. Then, for each $C > 0$, the resulting solution $Z(r)$ is smooth and differentiable in the interval $(0, \infty)$ with the following behaviour near the origin and infinity:

$$Z = \sigma_0 \left(\frac{1}{r} + \ldots + p(C)r^2 + O(r^3) \right) \qquad \text{as} \quad r \to 0,$$

$$Z = -\frac{2}{r^2} + \ldots + \frac{q(C)}{r^4} + O(\frac{ln(r)}{r^5}) \qquad \text{as} \quad r \to \infty. \tag{70}$$

Here dots denote terms specified entirely by the background solutions (12), (13), whereas the quantities $p(C)$ and $q(C)$ depend on the constant C as well. Using (16) we can return back to the tortoise coordinate ρ thus obtaining

$$Z = \frac{1}{\rho} + O(\rho) \ \text{ as } \rho \to 0, \quad \text{and} \ \ Z = -\frac{2}{\rho^2} + O(\frac{ln(\rho)}{\rho^3}) \ \text{ as } \rho \to \infty. \tag{71}$$

We therefore arrive at the one-parameter family of solutions $Z(\rho)$, whose members are distinguished by values of the constant $C > 0$ in (50), each solution satisfies boundary condition (71). Some of the typical solutions are presented in Fig.1. All these solutions have only one zero in the interval $(0, \infty)$ – this is because, in view of Eq.(46), the derivative Z' is always negative when Z vanishes. One should stress that such a behaviour of Z is common for all background regular equilibrium solutions.

Now, having obtained the function Z, we can determine the effective potential $U(\rho)$ in the Schrödinger equation (47). Notice that the first term in (48) contains an unbounded piece: $\sigma^2 N(3f^2 - 1)/r^2 \to 2/\rho^2$ when $\rho \to 0$. However, in account of condition (71), this divergency is exactly canceled by the second term in (48), such that the resulting potential turns out to be continuous in the whole semi-axis, $\rho \in [0, \infty)$, with the following behaviour near the origin and infinity:

$$U(\rho) = 6b(6b-1)\sigma_0^2 + O(\rho) \ \text{ as } \ \rho \to 0, \ \ U(\rho) = \frac{6}{\rho^2} + O(\frac{ln(\rho)}{\rho^3}) \ \text{ as } \ \rho \to \infty, \tag{72}$$

where b comes from Eq.(12). We therefore find a whole family of regular potentials. Each member of this family, $U(\rho)$, is specified by the corresponding solutions $Z(\rho)$ and satisfies boundary conditions (72). A typical example of such a potential is depicted in Fig.2.



Figure 1: Behavior of the function $Z(\rho)$ for $n = 1$ and $n = 2$ background BK solutions. Instead of Z, it is convenient to display Q such that $Z = \sigma Q/r$. For each n, a pair of solutions is shown corresponding to the two different choices of the integration constant $p(C)$ in Eq.(70) (the higher maximums correspond to $p(C) = 0$).



Figure 2: Odd-parity and even-parity potentials for the $n = 1$ background BK solution. The odd-parity potential is given by Eq.(48), the corresponding Z is shown in Fig.1 for $p(C) = 0$. The even-parity potential is defined by Eq.(29).

Thus the potential U in the Schrödinger equation (47) indeed can be chosen to be regular everywhere in the half-line $[0, \infty)$. However, U is not uniquely defined. Our aim now is to show that all potentials from the family specified above are equivalent as far as the bound state problem is concerned. To this end we study the properties of the transformations (45), (49).

Consider a solution of the initial system of equations (41), (42) corresponding to a bound state with $\omega^2 < 0$. Then, near the origin and infinity,

$$\xi \sim \rho \ , \quad \chi \sim \rho^2 \quad \text{as } \rho \to 0, \qquad\qquad \xi \sim \pm\frac{1}{\omega}\,\chi \sim exp(-|\omega|\rho) \quad \text{as} \ \ \rho \to \infty. \tag{73}$$

Applying transformation (45) with some particular $Z(\rho)$ and taking (71) into account one obtains the solution of Eq.(47) with the following behaviour:

$$\psi \sim \rho \ \text{ as } \ \rho \to 0, \qquad \psi \sim exp(-|\omega|\rho) \ \text{ as } \ \rho \to \infty, \tag{74}$$

which ensures normalizability of ψ

$$\langle \psi|\psi \rangle = \int_0^\infty \psi^2 \, d\rho < \infty; \tag{75}$$

so that ψ corresponds to the bound state solution of the Schrödinger equation (47) with the same $\omega^2 < 0$. On the other hand, starting from (74) and applying the inverse transformation (49), one arrives at (73). It is worth noting that these transformations uniquely fix the value $\psi(0) = 0$ ensuring essential self-adjointness of the Schrödinger operator $H = -d^2/d\rho^2 + U(\rho)$ (the Schrödinger equation by itself does not specify $\psi(0)$ because the potential is globally regular in the case).

Thus, for each particular U, transformations (45), (49) establish one-to-one correspondence between bound states with non-zero energy for the initial system of equations (41), (42) and those for the Schrödinger equation (47). Obviously, this means that the different choices of the potential U are equivalent. Precisely, the nature of this equivalency is explained by the following considerations. Let Z_1 and Z_2 be two different solutions of Eq.(46) (relating of course to the same background field configuration), and U_1, U_2 are the corresponding potentials. Then, given a solution of (47) for the potential U_1 and some $\omega^2 \neq 0$, ψ_1, one can construct a solution for the potential U_2 and *the same* ω^2 due to the following rule:

$$\psi_2 = \psi_1 + \frac{1}{\omega^2} f (Z_2 - Z_1) \left(f^3 Z_1 \psi_1 + f' \psi_1 - f \psi_1' \right). \tag{76}$$

Suppose that ψ_1 is a normalizable solution corresponding to a bound state of U_1, then it satisfies (74). From (70), (16) one concludes that

$$Z_1 - Z_2 \sim \rho^2 \ \text{ as } \ \rho \to 0, \qquad Z_1 - Z_2 \sim \frac{1}{\rho^4} \ \text{ as } \ \rho \to \infty, \tag{77}$$

which ensures that ψ_2 also satisfies (74), that is, it relates to a bound state of U_2 with the same eigenvalue $\omega^2 < 0$. Thus the bound state problem for Eqs.(41), (42) is equivalent to that for the Schrödinger equation (47) with any of the effective potentials from the family specified above.

7 Number of bound states

Our purpose now is to show that Eq.(47) admits a zero energy bound state. The simplest way to see this is to start from Eqs.(41), (42) which are known to possess the pure gauge solutions with zero energy. These solutions are given by Eq.(40) with the gauge parameter Ω satisfying Eq.(43). Then, substituting (40) in (45) one obtains a zero energy solution for Eq.(47), $\psi_0 = f\phi$, where $\phi = \Omega + Z\Omega'/\gamma^2$. Next, observing that the function ϕ obeys the following equation, $\phi' = f^2 Z\phi$, one can represent ψ_0 as

$$\psi_0 = f \exp\left(\int_{\rho_0}^{\rho} f^2 Z d\rho\right), \tag{78}$$

with some positive ρ_0; the validity of this solution can also be checked directly. The exponential factor in this expression is manifestly positive when $0 < \rho < \infty$ and, on account of the boundary conditions for Z, vanishes at $\rho = 0, \infty$, such that

$$\psi_0 \sim \rho \quad \text{as} \quad \rho \to 0, \qquad \psi_0 \sim \frac{1}{\rho^2} \quad \text{as} \quad \rho \to \infty, \tag{79}$$

which ensures normalizability of ψ_0. Thus ψ_0, being a normalizable solution of (47) and satisfying the right boundary condition at the origin, $\psi_0(0) = 0$, relates in fact to zero energy bound state. One should stress that this particular bound state occurs only for the Schrödinger equation (47) and not for the initial system of equations (41), (42). In fact, the equivalence between the initial system and the Schrödinger equation holds only for non-zero energy. When $\omega^2 = 0$, transformation (49) becomes ill-defined, and the equivalence is lost. Then, taking into account Eq.(69) one can see that the pure gauge solutions (40) by themselves can be neither normalizable nor even just regular both at the origin and infinity. However, their certain linear combination, ψ_0, can be normalizable. The idea is to use ψ_0 to establish the existence of *negative energy* bound states for the Schrödinger equation, and those are certainly equivalent to bound states appearing in the initial system.

To this end we observe that, owing to the oscillating factor f in (78), ψ_0 has exactly n nodes, which clearly proves that the Schrödinger equation admits n bound state solutions with negative energies. We therefore conclude that for the n-th regular BK solution there exist precisely n negative modes in the odd-parity spherical perturbation sector.

8 Black hole case

For black holes, the proof is essentially the same as that in the regular case. We start again form Eq.(43) and want to specify a particular solution for this equation, $\tilde{\Omega}(\rho)$, such that the corresponding $Z(\rho)$ is well behaved. The variable ρ now runs over the line $-\infty < \rho < \infty$, such that $\rho = -\infty$ (horizon) and $\rho = +\infty$ are regular singular points of the differential equation. Instead of ρ, it is convenient to use the independent variable x defined by (15), then Eq.(64) takes the form

$$x(1+x)^2\ \Omega''_{xx} + 2x(1+x)\left(1 - \left(f'_x\right)^2/r_h^2\right)\Omega'_x = 2f^2\frac{x}{N}\ \Omega. \tag{80}$$

In this equation, the coefficients are analytic functions in the vicinity of $x = 0$, because the background solutions (14) are analytic in some neighborhood of the horizon [10]. This ensures the existence of the fundamental system of solutions in the vicinity of $x = 0$. We choose the particular solution $\tilde{\Omega}$ whose regular integral vanishes at the horizon:

$$\tilde{\Omega}(x) = x(1 + X(x)), \quad \text{as} \quad x \to 0, \tag{81}$$

where $X(x)$ is given by a power series convergent in the vicinity of $x = 0$, $X(0) = 0$. Since the coefficients in (80) are at least continuous in the domain $x > 0$, the solution (81) admits at a smooth extension into this region. When x tends to infinity, the solution is given by (66) where, repeating the arguments used in Eqs.(67), (68), the coefficient before the growing part can be shown to be non-vanishing. From here one concludes that on the domain $0 < x < \infty$ there exists a smooth solution of (80) with the following behaviour near $x = 0, \infty$:

$$\tilde{\Omega} = x + O(x^2) \ \text{ as } x \to 0, \quad \text{ and } \ \tilde{\Omega} = const\ x + O(1) \ \text{ as } x \to \infty. \tag{82}$$

Next, using (16), (17) we express this solution in terms of ρ and substitute it into Eq.(50), where, in order to ensure the non-vanishing of the denominator, we choose the constant C such that

$$C < -\int_0^\infty f^2 \Lambda^2 d\rho; \tag{83}$$

the integral here exists since Eq.(82) ensures that $\Lambda \sim 1/\rho^2$ when $\rho \to \infty$. For each certain C we obtain from (50) a solution $Z(\rho)$ which is continuous and differentiable in $(-\infty, +\infty)$ with the following behaviour near the horizon and infinity:

$$Z = -\frac{1}{f_h^2 \rho} + O(\frac{1}{\rho^2}) \ \text{ as } \ \rho \to -\infty, \quad \text{ and } \ Z \to -\frac{2}{\rho^2} + O(\frac{ln(\rho)}{\rho^3}) \ \text{ as } \ \rho \to \infty. \tag{84}$$

The corresponding effective potential $U(\rho)$ is smooth and bounded. The equivalency of the different particular U can be checked with the use of (76), (84). Then, repeating the analysis performed above for the regular case, we arrive at the zero energy solution ψ_0 given by (78). In view of (84), this function satisfies the following boundary conditions:

$$\psi_0 \sim \frac{1}{\rho} \ \text{ as } \ \rho \to -\infty, \qquad \psi_0 \sim \frac{1}{\rho^2} \ \text{ as } \ \rho \to \infty \tag{85}$$

and is therefore normalizable on the whole line $(-\infty, +\infty)$. Thus in the black hole case there exists a zero energy bound state too. The function ψ_0 possesses n nodes, which shows the existence of n negative energy bound states. From here we conclude that for each (n, r_h) EYM black hole solution, $r_h \in (0, \infty)$, there exist precisely n negative modes in the odd-parity spherical perturbation sector. For the $n = 1$ EYM black holes, the negative modes energy eigenvalues for the both parity types are presented numerically in Tab.1.

Tab.1. The bound state energies for the $n = 1$ EYM black holes.

r_h	ω^2 (even-parity)	ω^2 (odd-parity)
0 (regular case)	–0.05249	–0.06190
0.1	–0.05026	–0.06182
0.5	–0.04021	–0.06000
1	–0.02683	–0.04926
5	–0.00212	–0.00298
10	–0.00054	–0.00075

Finally, we consider the case of the charged $SU(2) \times U(1)$ EYM black holes [18] which possess the additional $U(1)$ dyon type field. In this case, the only new thing appears – an extra term $q^2/2r^2$ in the right hand side of Eq.(9), where $q^2 = q_e^2 + q_m^2$, q_e and q_m being the dyon electric and magnetic charges; the odd-parity perturbation equations remain intact. The background solutions are qualitatively the same as in the $q = 0$ case, but the presence of the charge entails now that $r_h \geq q$. When $r_h > q$, all of the analysis performed above for the $q = 0$ case remains valid. The only difference arises in the extreme limit, $r_h = q$, when N has zero of second order at the horizon, and $f^2(r_h) = 1$. After all, the change of the boundary condition at the horizon results in the following change in the asymptotic behaviour for ψ_0 given by (78):

$$\psi_0 \sim \frac{1}{\rho^2} \quad \text{as} \quad \rho \to -\infty, \qquad \psi_0 \sim \frac{1}{\rho^2} \quad \text{as} \quad \rho \to \infty. \tag{86}$$

One can understand this as follows: for extreme black holes the geometry in the limit $\rho \to -\infty$ is known to be conformally flat. The dynamics of the conformally invariant Yang-Mills field then must be the same as that in the asymptotically flat region, $\rho \to \infty$ (remind that the odd-parity modes involve perturbations of the Yang-Mills field alone). Thus ψ_0 is normalizable in the case as well, and we therefore conclude that all charged $SU(2)\times U(1)$ black holes, including extreme solutions, also possess n odd-parity spherical negative modes.

Acknowledgments

I would like to thank Professor N.Straumann and also O.Brodbeck for discussions.

This work was supported by the Swiss National Science Foundation.

Appendix A Variation of the energy functional

In this Appendix second variation of the ADM mass functional presented in the main text is explicitly derived. The analysis is carried out for the regular BK solution, the generalization on the black hole case is straightforward.

Consider the barrier height functional defined by Eq.(30):

$$\varepsilon[f(r)] = \int_0^\infty (f'^2 + \frac{(f^2-1)^2}{2r^2}) exp(-2\int_r^\infty f'^2 \frac{dr}{r}) dr. \tag{A1}$$

Let $f(r)$ be a smooth function satisfying the following conditions

$$f'(r) = O(r) \quad \text{as } r \to 0; \qquad f'(r) = O(1/r^2) \quad \text{as } r \to \infty. \tag{A2}$$

Define for convenience two new functions

$$\sigma(r) = exp\{-2\int_r^\infty f'^2 \frac{dr}{r}\}, \quad m(r) = \frac{1}{\sigma(r)}\int_0^r (f'^2 + \frac{(f^2-1)^2}{r^2})\sigma dr, \tag{A3}$$

they satisfy the following boundary conditions

$$\sigma(\infty) = 1, \quad \sigma(0) \neq 0; \quad m(\infty) < \infty, \quad m(r) = O(r^3) \text{ as } r \to 0, \tag{A4}$$

and also, by definition, the following equations

$$\sigma' = 2\frac{f'^2}{r}\sigma, \qquad (m\sigma)' = (f'^2 + \frac{(f^2-1)^2}{2r^2})\sigma. \tag{A5}$$

Consider small variation

$$f(r) \ \to \ f(r) + \varphi(r), \tag{A6}$$

where

$$\varphi(0) = \varphi(\infty) = 0. \tag{A7}$$

To preserve the boundary conditions, the variation must also satisfy

$$\varphi'(r) = O(r) \text{ as } r \to 0, \quad \varphi'(r) = O(1/r^2) \text{ as } r \to \infty. \tag{A8}$$

Putting (A6) into (A1) and expanding the result over φ, one obtains

$$\varepsilon[f+\varphi] = \varepsilon[f] + \delta\varepsilon + \delta^2\varepsilon + \dots, \tag{A9}$$

where the first variation is

$$\delta\varepsilon = 2\int_0^\infty \{f'\varphi'\sigma + \frac{f(f^2-1)}{r^2}\sigma\varphi - 2I(f'^2 + \frac{(f^2-1)^2}{2r^2})\sigma\}dr, \tag{A10}$$

and the second variation is

$$\delta^2\varepsilon = \int_0^\infty \{\sigma\varphi^2 + \sigma\frac{3f^2-1}{r^2}\varphi^2 - 8I(f'\varphi' + \frac{f(f^2-1)}{r^2}\varphi)\sigma +$$

$$+ (8I^2 - 2J)(f'^2 + \frac{(f^2-1)^2}{2r^2})\sigma\}dr, \tag{A11}$$

dots in (A9) denote higher order terms, and the following new functions have been introduced:

$$I(r) = \int_r^\infty f'\varphi'\frac{dr}{r}, \quad J(r) = \int_r^\infty \varphi'^2\frac{dr}{r}. \tag{A12}$$

The boundary conditions (A2),(A8) imply that

$$I(0) < \infty, \ J(0) < \infty, \ I(\infty) = J(\infty) = 0. \tag{A13}$$

Consider the first variation (A10). Using (A5), represent $\delta\varepsilon$ as follows

$$\delta\varepsilon = 2\int_0^\infty \{f'\varphi'\sigma + \frac{f(f^2-1)}{r^2}\varphi\sigma - 2I(m\sigma)'\}dr. \tag{A14}$$

Integrating by parts one has

$$\delta\varepsilon = (2f'\sigma\varphi - 4Im\sigma)|_0^\infty + 2\int_0^\infty \{-(f'\sigma)'\varphi + \sigma\frac{f(f^2-1)}{r^2}\varphi + 2m\sigma I'\}dr. \tag{A15}$$

Note, that the boundary terms in this expression vanish. Finding I' from (A12) and integrating by parts once more, one finally arrives at

$$\delta\varepsilon = 2\int_0^\infty \{-((1-\frac{2m}{r})\sigma f')' + \frac{f(f^2-1)}{r^2}\sigma\}\varphi dr. \tag{A16}$$

One can see that the vanishing of the first variation implies the Yang-Mills equation

$$((1-\frac{2m}{r})\sigma f')' = \sigma\frac{f(f^2-1)}{r^2}. \tag{A17}$$

Assume now that the first variation vanishes and consider the second variation. Using (A5) one obtains

$$\delta^2\varepsilon = \int_0^\infty \{\sigma\varphi'^2 + \sigma\frac{3f^2-1}{r^2}\varphi^2 - 8I(f'\varphi' + \frac{f(f^2-1)}{r^2}\varphi)\sigma +$$

$$+\ 8I^2(m\sigma)' - 2J(m\sigma)'\}dr. \tag{A18}$$

Integrating the fourth term in the integrand by parts, and using Eq.(A12), one obtains

$$8\int_0^\infty I^2(m\sigma)'dr = 8I^2m\sigma\Big|_0^\infty - 16\int_0^\infty II'm\sigma dr = 16\int_0^\infty Im\sigma f'\varphi'\frac{dr}{r}, \tag{A19}$$

where the boundary terms vanish. Combining this result with the third term in Eq.(A18), one has

$$-8\int_0^\infty I\{(1-\frac{2m}{r})\sigma f'\varphi' + \frac{f(f^2-1)}{r^2}\sigma\varphi\}dr. \tag{A20}$$

Using (A17),(A12),(A5), represent this expression as follows

$$-8\int_0^\infty I((1-\frac{2m}{r})\sigma f'\varphi)'dr = -\ 8I(1-\frac{2m}{r})\sigma f'\varphi\Big|_0^\infty +$$

$$+\ 8\int_0^\infty I'(1-\frac{2m}{r})\sigma f'\varphi dr = -8\int_0^\infty(1-\frac{2m}{r})\sigma f'^2\varphi\varphi'\frac{dr}{r} =$$

$$= -4\int_0^\infty(1-\frac{2m}{r})\sigma'\varphi\varphi'dr, \tag{A21}$$

where the boundary terms vanish. The fifth term in (A18) yields

$$-2\int_0^\infty J(m\sigma)'dr = -2Jm\sigma\Big|_0^\infty + 2\int_0^\infty J'm\sigma dr = -2\int_0^\infty m\sigma\varphi'2\frac{dr}{r} =$$

$$= -2m\sigma\frac{\varphi}{r}\varphi'\Big|_0^\infty + 2\int_0^\infty(m\sigma\frac{\varphi'}{r})'\varphi dr = \int_0^\infty(\frac{2m}{r}\sigma\varphi')'\varphi dr. \tag{A22}$$

The first term in (A18) is

$$\int_0^\infty \sigma\varphi'^2 dr = -\int_0^\infty(\sigma\varphi')'dr, \tag{A23}$$

where the boundary terms are zero. Consider also the following expression

$$0 = \int_0^\infty((1-\frac{2m}{r})\sigma'\varphi^2)'dr. \tag{A24}$$

Adding the equations (A21)-(A24) and introducing also the tortoise coordinate, ρ,

$$\frac{dr}{d\rho} = \sigma(1 - \frac{2m}{r}), \tag{A25}$$

one finally arrives at

$$\delta^2\varepsilon = \int_0^\infty \varphi(-\frac{d^2}{d\rho^2} + V)\varphi \; d\rho, \tag{A26}$$

where

$$V = \sigma(1 - \frac{2m}{r})\{2(\sigma'(1 - \frac{2m}{r}))' + \frac{3f^2 - 1}{r^2}\sigma\}, \tag{A27}$$

which agrees with the potential in Eq.(29) provided that Eqs.(A5), (A17) are taken into account.

References

[1] R. Bartnik, J. McKinnon, Phys. Rev. Lett., **61** (1988) 141.

[2] M.S. Volkov, D.V. Gal'tsov, Pis'ma Zh. Eksp. Teor. Fiz., **50** (1989) 312
[JETP Lett., **50** (1990) 345]; Sov. J. Nucl.Phys. 51 (1990) 747;
H.P. Kunzle, A.K.M. Masood–ul–Alam, J. Math. Phys. **31** (1990) 928;
P. Bizon, Phys. Rev. Lett., **64** (1990) 2644.

[3] N. Straumann, Z.H. Zhou, Phys. Lett., **B 237** (1990) 353;
N. Straumann, Z.H. Zhou, Phys.Lett., **B 243** (1991) 53.
Z.H. Zhou, N. Straumann, Nucl. Phys., **B 369** (1991) 180;
Z.H. Zhou, Helv. Phys. Acta, **65** (1992) 767.

[4] D.V. Gal'tsov, M.S. Volkov, Phys. Lett., **B 273** (1991) 255;

[5] N.S.Manton, Phys.Rev., **D 28** (1983) 2019.

[6] F.R.Klinkhamer, N.S.Manton, Phys.Rev., **D 30** (1984) 2212.

[7] P.Forgacs, Z.Horath, Phys.Lett., **B 138** (1984) 397.

[8] J.A.Smoller, A.G.Wasserman, S.T.Yau, and J.B.McLeod,
Comm.Math.Phys. **143** (1991) 115.

[9] J.A.Smoller, A.G.Wasserman,Comm.Math.Phys. **151** (1993) 303;
ibid. **154** (1993) 377.

[10] P. Breitenlohner, P. Forgacs, D. Maison, Comm. Math. Phys. **163** (1994) 141.

[11] J.Burzlaff, Nucl.Phys., **B 233** (1984) 262.

[12] M.S. Volkov, Phys. Lett., **B 334** (1994) 40.

[13] M.S. Volkov, Phys. Lett., **B 328** (1994) 89.

[14] M.S.Volkov, D.V.Gal'tsov, Phys.Lett., **B 341** (1995) 279.

[15] O. Brodbeck, N. Straumann, Phys. Lett., **B 324** (1994) 309.

[16] G.Lavrelashvili, D.Maison., Phys.Lett., **343** (1995) 214.

[17] M.S.Volkov, O.Brodbeck, G.Lavrelashvili, N.Straumann, Phys.Lett., **B 349** (1995) 438.

[18] D.V. Gal'tsov, M.S. Volkov, Phys. Lett., **B 274** (1992) 173;

[19] E.T.Whittaker, G.N.Watson, *A Course of Modern Analysis*, Cambridge University Press (1927).